

ANTI-DEPRESSANTS

MEDICATION(S) SUBJECT TO STEP THERAPY

TRINTELLIX, VIIBRYD, VIIBRYD STARTER PACK

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, Paxil suspension, sertraline, venlafaxine.

ANTI-SEIZURE DRUGS

MEDICATION(S) SUBJECT TO STEP THERAPY

APTOM, FYCOMPA, VIMPAT 10 MG/ML SOLUTION, VIMPAT 100 MG TAB, VIMPAT 150 MG TAB, VIMPAT 200 MG TAB, VIMPAT 50 MG TAB

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): carbamazepine, clonazepam, divalproex, felbamate, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, tiagabine, topiramate, zonisamide.

BRIVARACETAM (BRIVIACT)

MEDICATION(S) SUBJECT TO STEP THERAPY

BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB

CRITERIA

The following drug need to be used first (step 1 drugs): levetiracetam.

DESVENLAFAXINE ER TABLET

MEDICATION(S) SUBJECT TO STEP THERAPY

DESVENLAFAXINE ER

CRITERIA

One of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er (generic Pristiq), duloxetine, fluoxetine, fluvoxamine, paroxetine, Paxil oral suspension, sertraline, venlafaxine, escitalopram

FLUNISOLIDE NASAL SPRAY (FLUNISOLIDE)

MEDICATION(S) SUBJECT TO STEP THERAPY

FLUNISOLIDE

CRITERIA

Following drug needs to be used first (step 1 drug): fluticasone propionate nasal spray.

NON-PREFERRED LOW POTENCY TOPICAL STEROIDS

MEDICATION(S) SUBJECT TO STEP THERAPY

DESONIDE 0.05 % CREAM, DESONIDE 0.05 % OINTMENT

CRITERIA

Trial of two preferred topical steroids from the low potency group (step 1 drugs): alclometasone dipropionate (cream, ointment), hydrocortisone (cream, ointment, lotion), betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution), fluocinolone (cream, ointment), fluticasone (cream, ointment), triamcinolone (cream, lotion, ointment).

NON-PREFERRED MEDIUM POTENCY TOPICAL STEROIDS

MEDICATION(S) SUBJECT TO STEP THERAPY

HYDROCORTISONE BUTYRATE 0.1 % OINTMENT, HYDROCORTISONE BUTYRATE 0.1 % SOLUTION,
HYDROCORTISONE VALERATE 0.2 % CREAM

CRITERIA

Trial of two preferred topical steroids from the medium potency group (step 1 drugs):
betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution),
fluocinolone (cream, ointment), fluticasone (cream, ointment), triamcinolone (cream, lotion,
ointment).

OVERACTIVE BLADDER AGENTS

MEDICATION(S) SUBJECT TO STEP THERAPY

TOLTERODINE TARTRATE, TOLTERODINE TARTRATE ER

CRITERIA

One of the following drugs need to be used first (step 1 drugs): oxybutynin, trospium.

RUFINAMIDE (BANZEL)

MEDICATION(S) SUBJECT TO STEP THERAPY

RUFINAMIDE

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): clonazepam, divalproex sodium, felbamate, lamotrigine, topiramate, valproic acid.

TACROLIMUS OINTMENT (PROTOPIC OINTMENT)

MEDICATION(S) SUBJECT TO STEP THERAPY

TACROLIMUS 0.03 % OINTMENT, TACROLIMUS 0.1 % OINTMENT

CRITERIA

One of the following topical steroids need to be used first (step 1 drugs): betamethasone dipropionate (cream, ointment, lotion), betamethasone augmented (gel, lotion, cream, ointment), betamethasone valerate (cream, ointment, lotion), clobetasol (cream, ointment, gel, solution), clobetasol emollient cream, fluocinolone (cream, ointment), fluocinonide (gel, ointment, solution), fluocinonide emollient cream, fluticasone (cream, ointment), halobetasol (cream, ointment), hydrocortisone valerate (cream, ointment), mometasone (cream, ointment, solution), triamcinolone (cream, lotion, ointment).