

## **ANTI-DEPRESSANTS**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

TRINTELLIX, VILAZODONE HCL

### **CRITERIA**

Two of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine.

## **BIMATOPROST EYE DROPS (BIMATOPROST)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

BIMATOPROST

### **CRITERIA**

The following drug needs to be used first (step 1 drug): latanoprost eye drops.

## **BRIVARACETAM (BRIVIACT)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB

### **CRITERIA**

The following drug need to be used first (step 1 drugs): levetiracetam.

## **CARVEDILOL EXTENDED RELEASE (COREG CR)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

CARVEDILOL PHOSPHATE ER

### **CRITERIA**

The following drug needs to be used first (step 1 drugs): carvedilol immediate release.

## **DONEPEZIL 23MG (ARICEPT)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

DONEPEZIL HCL 23 MG TAB

### **CRITERIA**

One of the following drugs need to be used first (step 1 drugs): donepezil 5mg, donepezil 10mg.

## **FEBUXOSTAT (ULORIC)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

FEBUXOSTAT

### **CRITERIA**

The following drug needs to be used first (step 1 drugs): allopurinol.

## **NON-PREFERRED MEDIUM POTENCY TOPICAL STEROIDS**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

HYDROCORTISONE BUTYRATE 0.1 % OINTMENT

### **CRITERIA**

Trial of two preferred topical steroids from the medium potency group (step 1 drugs):  
betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution),  
fluocinolone (cream, ointment), fluticasone (cream, ointment), triamcinolone (cream, lotion,  
ointment).

## OVERACTIVE BLADDER AGENTS

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

TOLTERODINE TARTRATE, TOLTERODINE TARTRATE ER

### **CRITERIA**

One of the following drugs need to be used first (step 1 drugs): oxybutynin, trospium, solifenacin

## **RUFINAMIDE (BANZEL)**

---

### **MEDICATION(S) SUBJECT TO STEP THERAPY**

RUFINAMIDE

### **CRITERIA**

Two of the following drugs need to be used first (step 1 drugs): clonazepam, divalproex sodium, felbamate, lamotrigine, topiramate.