

## ANTI-DEPRESSANTS

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

TRINTELLIX, VIIBRYD, VIIBRYD STARTER PACK

### CRITERIA

Two of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, Paxil suspension, sertraline, venlafaxine.

## ANTI-SEIZURE DRUGS

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

APTOM, FYCOMPA, VIMPAT 10 MG/ML SOLUTION, VIMPAT 100 MG TAB, VIMPAT 150 MG TAB, VIMPAT 200 MG TAB, VIMPAT 50 MG TAB

### CRITERIA

Two of the following drugs need to be used first (step 1 drugs): carbamazepine, clonazepam, divalproex, felbamate, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, tiagabine, topiramate, zonisamide.

## BRIVARACETAM (BRIVIACT)

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB

### CRITERIA

The following drug need to be used first (step 1 drugs): levetiracetam.

## FENOFIBRATE MICRONIZED (ANTARA)

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

FENOFIBRATE MICRONIZED 130 MG CAP, FENOFIBRATE MICRONIZED 43 MG CAP

### CRITERIA

One of the following drugs need to be used first (step 1 drugs): fenofibrate

## FLUNISOLIDE NASAL SPRAY (FLUNISOLIDE)

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

FLUNISOLIDE

### CRITERIA

Following drug needs to be used first (step 1 drug): fluticasone propionate nasal spray.

## NON-PREFERRED MEDIUM POTENCY TOPICAL STEROIDS

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

HYDROCORTISONE BUTYRATE 0.1 % OINTMENT, HYDROCORTISONE BUTYRATE 0.1 % SOLUTION,  
HYDROCORTISONE VALERATE 0.2 % CREAM

### CRITERIA

Trial of two preferred topical steroids from the medium potency group (step 1 drugs):  
betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution),  
prednicarbate ointment, fluocinolone (cream, ointment), fluticasone (cream, ointment),  
triamcinolone (cream, lotion, ointment).

## RUFINAMIDE (BANZEL)

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

RUFINAMIDE

### CRITERIA

Two of the following drugs need to be used first (step 1 drugs): clonazepam, divalproex sodium, felbamate, lamotrigine, topiramate, valproic acid.

## TACROLIMUS OINTMENT (PROTOPIC OINTMENT)

---

### MEDICATION(S) SUBJECT TO STEP THERAPY

TACROLIMUS 0.03 % OINTMENT, TACROLIMUS 0.1 % OINTMENT

### CRITERIA

One of the following topical steroids need to be used first (step 1 drugs): betamethasone dipropionate (cream, ointment, lotion), betamethasone augmented (gel, lotion, cream, ointment), betamethasone valerate (cream, ointment, lotion), clobetasol (cream, ointment, gel, solution), clobetasol emollient cream, fluocinolone (cream, ointment), fluocinonide (gel, ointment, solution), fluocinonide emollient cream, fluticasone (cream, ointment), halobetasol (cream, ointment), hydrocortisone valerate (cream, ointment), mometasone (cream, ointment, solution), prednicarbate ointment, triamcinolone (cream, lotion, ointment).