

ADAPALENE-BENZOYL PEROXIDE (EPIDUO, EPIDUO FORTE)

MEDICATION(S) SUBJECT TO STEP THERAPY

ADAPALENE-BENZOYL PEROXIDE 0.1-2.5 % GEL, ADAPALENE-BENZOYL PEROXIDE 0.3-2.5 % GEL

CRITERIA

One of the following topical drugs need to be used first (step 1 drugs): adapalene or benzoyl peroxide-containing product.

ANTI-DEPRESSANTS

MEDICATION(S) SUBJECT TO STEP THERAPY

TRINTELLIX, VIIBRYD STARTER PACK, VILAZODONE HCL

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine.

BIMATOPROST EYE DROPS (BIMATOPROST)

MEDICATION(S) SUBJECT TO STEP THERAPY

BIMATOPROST

CRITERIA

The following drug needs to be used first (step 1 drug): latanoprost eye drops.

BRIVARACETAM (BRIVIACT)

MEDICATION(S) SUBJECT TO STEP THERAPY

BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB

CRITERIA

The following drug need to be used first (step 1 drugs): levetiracetam.

CARVEDILOL EXTENDED RELEASE (COREG CR)

MEDICATION(S) SUBJECT TO STEP THERAPY

CARVEDILOL PHOSPHATE ER

CRITERIA

The following drug needs to be used first (step 1 drugs): carvedilol immediate release.

CLINDAMYCIN/BENZOYL PEROXIDE GEL WITH PUMP

MEDICATION(S) SUBJECT TO STEP THERAPY

CLINDAMYCIN PHOS-BENZOYL PEROX 1.2-2.5 % GEL

CRITERIA

One of the following drugs need to be used first (step 1 drugs): clindamycin 1%/benzoyl peroxide 5% (generic BENZACLIN), clindamycin 1.2%/benzoyl peroxide 5% (generic DUAC).

CLINDAMYCIN/TRETINOIN GEL (VELTIN)

MEDICATION(S) SUBJECT TO STEP THERAPY

CLINDAMYCIN-TRETINOIN

CRITERIA

Both of the following drugs need to be used first (step 1 drugs): topical clindamycin, topical tretinoin.

DES Loratadine ODT (Clarinet)

MEDICATION(S) SUBJECT TO STEP THERAPY

DES Loratadine 2.5 MG TAB DISP, DES Loratadine 5 MG TAB DISP

CRITERIA

The following drug needs to be used first (step 1 drugs): desloratadine 5mg (non-dissolving) tablet.

DONEPEZIL 23MG (ARICEPT)

MEDICATION(S) SUBJECT TO STEP THERAPY

DONEPEZIL HCL 23 MG TAB

CRITERIA

One of the following drugs need to be used first (step 1 drugs): donepezil 5mg, donepezil 10mg.

FEBUXOSTAT (ULORIC)

MEDICATION(S) SUBJECT TO STEP THERAPY

FEBUXOSTAT

CRITERIA

The following drug needs to be used first (step 1 drugs): allopurinol.

FLUVOXAMINE EXTENDED RELEASE

MEDICATION(S) SUBJECT TO STEP THERAPY

FLUVOXAMINE MALEATE ER

CRITERIA

The following drug needs to be used first (step 1 drugs): fluvoxamine immediate release.

KETOCONAZOLE FOAM (EXTINA)

MEDICATION(S) SUBJECT TO STEP THERAPY

KETOCONAZOLE 2 % FOAM, KETODAN 2 % FOAM

CRITERIA

One of the following drugs need to be used first (step 1 drugs): ketoconazole 2% cream, ketoconazole 2% shampoo.

LAMOTRIGINE EXTENDED RELEASE (LAMICTAL XR)

MEDICATION(S) SUBJECT TO STEP THERAPY

LAMOTRIGINE ER

CRITERIA

The following drug needs to be used first (step 1 drugs): lamotrigine immediate release.

MOMETASONE FUROATE NASAL SPRAY (NASONEX)

MEDICATION(S) SUBJECT TO STEP THERAPY

MOMETASONE FUROATE 50 MCG/ACT SUSPENSION

CRITERIA

One of the following drugs need to be used first (step 1 drugs): flunisolide nasal spray, fluticasone nasal spray.

NON-PREFERRED HIGH POTENCY TOPICAL STEROIDS

MEDICATION(S) SUBJECT TO STEP THERAPY

DESOXIMETASONE 0.25 % LIQUID, FLUOCINONIDE 0.1 % CREAM

CRITERIA

Trial of two preferred topical steroids from the high potency group (step 1 drugs): betamethasone dipropionate (cream, ointment, lotion), fluocinonide (cream, gel, ointment, solution), fluocinonide emollient cream, triamcinolone 0.5% (cream, ointment).

NON-PREFERRED IMMEDIATE-RELEASE ADHD DRUGS

MEDICATION(S) SUBJECT TO STEP THERAPY

AMPHETAMINE SULFATE

CRITERIA

One of the following drugs need to be used first (step 1 drugs):

amphetamine/dexamphetamine (generic ADDERALL), dexamethylphenidate (generic FOCALIN), methylphenidate (generic METHYLIN), methylphenidate (generic RITALIN).

NON-PREFERRED MEDIUM POTENCY TOPICAL STEROIDS

MEDICATION(S) SUBJECT TO STEP THERAPY

DESOXIMETASONE 0.05 % GEL, DESOXIMETASONE 0.05 % OINTMENT, HYDROCORTISONE BUTYRATE 0.1 % CREAM, HYDROCORTISONE BUTYRATE 0.1 % OINTMENT, HYDROCORTISONE BUTYRATE 0.1 % SOLUTION, HYDROCORTISONE VALERATE 0.2 % CREAM

CRITERIA

Trial of two preferred topical steroids from the medium potency group (step 1 drugs): betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution), fluocinolone (cream, ointment), fluticasone (cream, ointment), triamcinolone (cream, lotion, ointment).

NON-PREFERRED MESALAMINE PRODUCT

MEDICATION(S) SUBJECT TO STEP THERAPY

MESALAMINE 400 MG CAP DR, MESALAMINE 800 MG TAB DR, MESALAMINE ER 500 MG CAP ER

CRITERIA

The following drugs need to be used first (step 1 drugs): balsalazide 750 capsule AND either mesalamine dr 1.2-gram tablet or mesalamine 0.375 gram er capsule.

NON-PREFERRED TOPICAL ANTI-ANTIFUNGAL

MEDICATION(S) SUBJECT TO STEP THERAPY

LULICONAZOLE, NAFTIFINE HCL 1 % CREAM, NAFTIFINE HCL 2 % CREAM, OXICONAZOLE NITRATE

CRITERIA

One of the following drugs need to be used first (step 1 drugs): ciclopirox (cream, gel, suspension, shampoo), clotrimazole (cream, solution), clotrimazole/betamethasone (cream, lotion), econazole nitrate cream, ketoconazole (cream, shampoo), nystatin (cream, ointment, powder, tablet), nystatin-triamcinolone (cream, ointment).

OVERACTIVE BLADDER AGENTS

MEDICATION(S) SUBJECT TO STEP THERAPY

DARIFENACIN HYDROBROMIDE ER, TOLTERODINE TARTRATE, TOLTERODINE TARTRATE ER

CRITERIA

One of the following drugs need to be used first (step 1 drugs): oxybutynin, trospium, solifenacin

RUFINAMIDE (BANZEL)

MEDICATION(S) SUBJECT TO STEP THERAPY

RUFINAMIDE

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): clonazepam, divalproex sodium, felbamate, lamotrigine, topiramate.

TAFLUPROST EYE DROPS (ZIOPTAN)

MEDICATION(S) SUBJECT TO STEP THERAPY

TAFLUPROST (PF)

CRITERIA

The following drug needs to be used first (step 1 drug): latanoprost eye drops

TRIAMTERENE (DYRENIUM)

MEDICATION(S) SUBJECT TO STEP THERAPY

TRIAMTERENE

CRITERIA

One of the following drugs need to be used first (step 1 drugs): amiloride, amiloride hctz, spironolactone, spironolactone hctz, or triamterene/hctz.