

ANTI-DEPRESSANTS

MEDICATION(S) SUBJECT TO STEP THERAPY

TRINTELLIX, VIIBRYD STARTER PACK, VILAZODONE HCL

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): bupropion, citalopram, desvenlafaxine succinate er, duloxetine, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline, venlafaxine.

BIMATOPROST EYE DROPS (BIMATOPROST)

MEDICATION(S) SUBJECT TO STEP THERAPY

BIMATOPROST

CRITERIA

The following drug needs to be used first (step 1 drug): latanoprost eye drops.

BRIVARACETAM (BRIVIACT)

MEDICATION(S) SUBJECT TO STEP THERAPY

BRIVIACT 10 MG TAB, BRIVIACT 10 MG/ML SOLUTION, BRIVIACT 100 MG TAB, BRIVIACT 25 MG TAB, BRIVIACT 50 MG TAB, BRIVIACT 75 MG TAB

CRITERIA

The following drug need to be used first (step 1 drugs): levetiracetam.

CARVEDILOL EXTENDED RELEASE (COREG CR)

MEDICATION(S) SUBJECT TO STEP THERAPY

CARVEDILOL PHOSPHATE ER

CRITERIA

The following drug needs to be used first (step 1 drugs): carvedilol immediate release.

DEXLANSOPRAZOLE (DEXILANT)

MEDICATION(S) SUBJECT TO STEP THERAPY

DEXLANSOPRAZOLE

CRITERIA

One of the following drugs need to be used first (step 1 drugs): lansoprazole, omeprazole, pantoprazole, rabeprazole.

DONEPEZIL 23MG (ARICEPT)

MEDICATION(S) SUBJECT TO STEP THERAPY

DONEPEZIL HCL 23 MG TAB

CRITERIA

One of the following drugs need to be used first (step 1 drugs): donepezil 5mg, donepezil 10mg.

FEBUXOSTAT (ULORIC)

MEDICATION(S) SUBJECT TO STEP THERAPY

FEBUXOSTAT

CRITERIA

The following drug needs to be used first (step 1 drugs): allopurinol.

NON-PREFERRED MEDIUM POTENCY TOPICAL STEROIDS

MEDICATION(S) SUBJECT TO STEP THERAPY

HYDROCORTISONE BUTYRATE 0.1 % OINTMENT, HYDROCORTISONE VALERATE 0.2 % CREAM

CRITERIA

Trial of two preferred topical steroids from the medium potency group (step 1 drugs): betamethasone valerate (cream, ointment, lotion), mometasone (cream, ointment, solution), fluocinolone (cream, ointment), fluticasone (cream, ointment), triamcinolone (cream, lotion, ointment).

OVERACTIVE BLADDER AGENTS

MEDICATION(S) SUBJECT TO STEP THERAPY

TOLTERODINE TARTRATE, TOLTERODINE TARTRATE ER

CRITERIA

One of the following drugs need to be used first (step 1 drugs): oxybutynin, trospium, solifenacin

RUFINAMIDE (BANZEL)

MEDICATION(S) SUBJECT TO STEP THERAPY

RUFINAMIDE

CRITERIA

Two of the following drugs need to be used first (step 1 drugs): clonazepam, divalproex sodium, felbamate, lamotrigine, topiramate.